

Fibonacci
(Leonardo Pisano)
1170-1240?
Statue in Pisa Italy



FIBONACCI NUMBERS
GOLDEN RATIO,
RECURRENCES

Lecture 27
CS2110 – Fall 2018

Announcements

A7: NO LATE DAYS. No need to put in time and comments. We have to grade quickly. No regrade requests for A7. Grade based only on your score on a bunch of sewer systems.

Please check submission guidelines carefully. Every mistake you make in submitting A7 slows down grading of A7 and consequent delay of publishing tentative course grades.

Announcements

Final is optional! As soon as we grade A7 and get it into the CMS, we determine tentative course grades.

You will complete “assignment” [Accept course grade?](#) on the CMS by Wednesday night.

If you accept it, that IS your grade. It won't change.

Don't accept it? Take final. Can lower and well as raise grade.

More past finals are now on [Exams page of course website](#). Not all answers yet.

Announcements

Course evaluation: Completing it is part of your course assignment. Worth 1% of grade.

Must be completed by Saturday night. 1 DEC

We then get a file that says who completed the evaluation.

We do not see your evaluations until after we submit grades to the Cornell system.

We never see names associated with evaluations.

Fibonacci function

$\text{fib}(0) = 0$

$\text{fib}(1) = 1$

$\text{fib}(n) = \text{fib}(n-1) + \text{fib}(n-2)$ for $n \geq 2$

0, 1, 1, 2, 3, 5, 8, 13, 21, ...

In his book in 1202 titled *Liber Abaci*

Has nothing to do with the famous pianist Liberaci

But sequence described much earlier in India:

Virahanka 600–800
Gopala before 1135
Hemacandra about 1150

The so-called Fibonacci numbers in ancient and medieval India.
[Parmanad Singh, 1985 pdf on course website](#)

Fibonacci function (year 1202)

$\text{fib}(0) = 0$

$\text{fib}(1) = 1$

$\text{fib}(n) = \text{fib}(n-1) + \text{fib}(n-2)$ for $n \geq 2$

```
/** Return fib(n). Precondition: n ≥ 0.*/
public static int f(int n) {
    if (n <= 1) return n;
    return f(n-1) + f(n-2);
}
```

0, 1, 1, 2, 3, 5, 8, 13, 21, 34, 55

We'll see that this is a lousy way to compute $f(n)$

Golden ratio $\Phi = (1 + \sqrt{5})/2 = 1.61803398\dots$

7

Divide a line into two parts:
 Call long part **a** and short part **b**



$(a + b) / a = a / b$

Solution is the golden ratio, Φ

See webpage:
<http://www.mathsisfun.com/numbers/golden-ratio.html>

$\Phi = (1 + \sqrt{5})/2 = 1.61803398\dots$

8

0, 1, 1, 2, 3, 5, 8, 13, 21, 34, 55

$fib(n) / fib(n-1)$ is close to Φ .
 So $\Phi * fib(n-1)$ is close to $fib(n)$
 Use formula to calculate $fib(n)$ from $fib(n-1)$

In fact,

limit $f(n)/fib(n-1) = \Phi$
 $n \rightarrow \infty$

a/b
8/5 = 1.6
13/8 = 1.625...
21/13 = 1.615...
34/21 = 1.619...
55/34 = 1.617...

Golden ratio and Fibonacci numbers: inextricably linked

Golden ratio $\Phi = (1 + \sqrt{5})/2 = 1.61803398\dots$

9

Find the golden ratio when we divide a line into two parts a and b such that

$(a + b) / a = a / b = \Phi$

Golden rectangle



a/b
8/5 = 1.6
13/8 = 1.625...
21/13 = 1.615...
34/21 = 1.619...
55/34 = 1.617...

For successive Fibonacci numbers a, b, a/b is close to Φ but not quite it Φ . 0, 1, 1, 2, 3, 5, 8, 13, 21, 34, 55, ...

Fibonacci, golden ratio, golden angle

10

0, 1, 1, 2, 3, 5, 8, 13, 21, 34, 55

limit $f(n)/fib(n-1) = \text{golden ratio} = 1.6180339887\dots$
 $n \rightarrow \infty$


360/1.6180339887... = 222.492235...

360 - 222.492235... = 137.5077 golden angle


Fibonacci function (year 1202)

11

Downloaded from wikipedia




Golden rectangle
 Fibonacci tiling



Fibonacci spiral

0, 1, 1, 2, 3, 5, 8, 13, 21, 34 ...



Drawing a golden rectangle with ruler and compass

13

golden rectangle

hypotenuse: $\sqrt{(1*1 + (\frac{1}{2})(\frac{1}{2}))} = \sqrt{(5/4)}$

How to draw a golden rectangle

fibonacci and bees

14

Male bee has only a mother
Female bee has mother and father

The number of ancestors at any level is a Fibonacci number

MB: male bee, FB: female bee

Fibonacci in Pascal's Triangle

15

Caching

31

As values of $f(n)$ are calculated, save them in an ArrayList. Call it a **cache**.

When asked to calculate $f(n)$ see if it is in the cache. If yes, just return the cached value. If no, calculate $f(n)$, add it to the cache, and return it.

Must be done in such a way that if $f(n)$ is about to be cached, $f(0), f(1), \dots, f(n-1)$ are already cached.

Caching

32

/** For $0 \leq n < \text{cache.size}$, $\text{fib}(n)$ is $\text{cache}[n]$
***** If $\text{fibCached}(k)$ has been called, its result is in $\text{cache}[k]$ ***/**
 public static ArrayList<Integer> cache= new ArrayList<>();

/** Return fibonacci(n). Pre: $n \geq 0$. Use the cache. ***/**
 public static int fibCached(int n) {
 if (n < cache.size()) return cache.get(n);
 if (n == 0) { cache.add(0); return 0; }
 if (n == 1) { cache.add(1); return 1; }

 int ans= fibCached(n-2) + fibCached(n-1);
 cache.add(ans);
 return ans;
 }

Linear algorithm to calculate fib(n)

33

/** Return fib(n), for $n \geq 0$. ***/**
 public static int f(int n) {
 if (n <= 1) return 1;
 int p=0; int c= 1; int i=2;
 // invariant: p = fib(i-2) and c = fib(i-1)
 while (i < n) {
 int fibi= c + p; p= c; c= fibi;
 i= i+1;
 }
 return c + p;
 }

Logarithmic algorithm!

34

$$\begin{aligned}
 f_0 &= 0 \\
 f_1 &= 1 \\
 f_{n+2} &= f_{n+1} + f_n
 \end{aligned}
 \quad
 \begin{pmatrix} 0 & 1 \\ 1 & 1 \end{pmatrix}
 \begin{pmatrix} f_n \\ f_{n+1} \end{pmatrix}
 =
 \begin{pmatrix} f_{n+1} \\ f_{n+2} \end{pmatrix}$$

$$\begin{pmatrix} 0 & 1 \\ 1 & 1 \end{pmatrix}
 \begin{pmatrix} 0 & 1 \\ 1 & 1 \end{pmatrix}
 \begin{pmatrix} f_n \\ f_{n+1} \end{pmatrix}
 =
 \begin{pmatrix} 0 & 1 \\ 1 & 1 \end{pmatrix}
 \begin{pmatrix} f_{n+1} \\ f_{n+2} \end{pmatrix}
 =
 \begin{pmatrix} f_{n+2} \\ f_{n+3} \end{pmatrix}$$

$$\begin{pmatrix} 0 & 1 \\ 1 & 1 \end{pmatrix}^k
 \begin{pmatrix} f_n \\ f_{n+1} \end{pmatrix}
 =
 \begin{pmatrix} f_{n+k} \\ f_{n+k+1} \end{pmatrix}$$

Logarithmic algorithm!

35

$$\begin{aligned}
 f_0 &= 0 \\
 f_1 &= 1 \\
 f_{n+2} &= f_{n+1} + f_n
 \end{aligned}
 \quad
 \begin{pmatrix} 0 & 1 \\ 1 & 1 \end{pmatrix}^k
 \begin{pmatrix} f_n \\ f_{n+1} \end{pmatrix}
 =
 \begin{pmatrix} f_{n+k} \\ f_{n+k+1} \end{pmatrix}$$

$$\begin{pmatrix} 0 & 1 \\ 1 & 1 \end{pmatrix}^k
 \begin{pmatrix} f_0 \\ f_1 \end{pmatrix}
 =
 \begin{pmatrix} f_k \\ f_{k+1} \end{pmatrix}$$

You know a logarithmic algorithm for exponentiation —recursive and iterative versions

Gries and Levin
 Computing a Fibonacci number in log time.
 IPL 2 (October 1980), 68-69.

Another log algorithm!

36

Define $\phi = (1 + \sqrt{5}) / 2$ $\phi' = (1 - \sqrt{5}) / 2$

The golden ratio again.

Prove by induction on n that

$$f_n = (\phi^n - \phi'^n) / \sqrt{5}$$